

On a Cuntz-Krieger functor

Igor V. Nikolaev *

Abstract

We construct a covariant functor from the topological torus bundles to the so-called Cuntz-Krieger algebras; the functor maps homeomorphic bundles into the stably isomorphic Cuntz-Krieger algebras. It is shown, that the K -theory of the Cuntz-Krieger algebra encodes torsion of the first homology group of the bundle. We illustrate the result by examples.

Key words and phrases: torus bundle, Cuntz-Krieger algebra

AMS (MOS) Subj. Class.: 46L; 57M

1 Introduction

A. The Cuntz-Krieger algebras. Recall, that the Cuntz-Krieger algebra, $\mathcal{O}_A$, is a C^* -algebra generated by partial isometries $s_1, \dots, s_n$; they act on the Hilbert space in such a way, that their support projections $Q_i = s_i^* s_i$ and their range projections $P_i = s_i s_i^*$ are orthogonal and satisfy the relations

$$Q_i = \sum_{j=i}^n a_{ij} P_j, \quad (1)$$

for an $n \times n$ matrix $A = (a_{ij})$ consisting of 0's and 1's [1]. The notion is extendable to the matrices A with the non-negative integer entries *ibid.*, *Remark 2.16*. It is known, that the C^* -algebra $\mathcal{O}_A$ is simple, whenever matrix A is irreducible; the latter means that certain power of A is a strictly

*Partially supported by NSERC.

positive integer matrix. If $\mathcal{K}$ is the C^* -algebra of compact operators on a Hilbert space, then the Cuntz-Krieger algebras $\mathcal{O}_A, \mathcal{O}_{A'}$ are said to be *stably isomorphic*, if $\mathcal{O}_A \otimes \mathcal{K} \cong \mathcal{O}_{A'} \otimes \mathcal{K}$, where $\cong$ is an isomorphism of the C^* -algebras. The K -theory of $\mathcal{O}_A$ was established in [1]; it was shown, that $K_0(\mathcal{O}_A) \cong \mathbb{Z}^n / (I - A^t)\mathbb{Z}^n$ and $K_1(\mathcal{O}_A) = \text{Ker} (I - A^t)$, where A^t is a transpose of the matrix A . It is easy to see, that if $\det (I - A^t) \neq 0$, then $K_0(\mathcal{O}_A)$ is a finite abelian group and $K_1(\mathcal{O}_A) = 0$; the two groups are invariants of the stable isomorphism.

B. The torus bundles. Let T^n be a torus of dimension $n \geq 1$. Recall, that *torus bundle* is an $(n + 1)$ -dimensional manifold

$$M_\alpha = \{T^n \times [0, 1] \mid (T^n, 0) = (\alpha(T^n), 1)\}, \quad (2)$$

where $\alpha : T^n \rightarrow T^n$ is an automorphism of T^n . It is known, that bundles M_α and $M_{\alpha'}$ are homeomorphic, whenever the automorphisms α and α' are conjugate, i.e. $\alpha' = \beta \circ \alpha \circ \beta^{-1}$ for an automorphism $\beta : T^n \rightarrow T^n$. Let $H_1(T^n; \mathbb{Z}) \cong \mathbb{Z}^n$ be the first homology of torus; consider the group $\text{Aut} (T^n)$ of (homotopy classes of) automorphisms of T^n . Any $\alpha \in \text{Aut} (T^n)$ induces a linear transformation of $H_1(T^n; \mathbb{Z})$, given by an invertible $n \times n$ matrix A with the integer entries; conversely, each $A \in GL_n(\mathbb{Z})$ defines an automorphism $\alpha : T^n \rightarrow T^n$. In this matrix representation, the conjugate automorphisms α and α' define similar matrices $A, A' \in GL_n(\mathbb{Z})$, i.e. such that $A' = BAB^{-1}$ for a matrix $B \in GL_n(\mathbb{Z})$. Each class of matrices, similar to a matrix $A \in GL_n(\mathbb{Z})$ and such that $\text{tr} (A) \geq 0$ ($\text{tr} (A) \leq 0$), contains a matrix with only the non-negative (non-positive) entries. We always assume, that our bundle M_α is given by a non-negative matrix A ; the matrices with $\text{tr} (A) \leq 0$ can be reduced to this case by switching the sign (from negative to positive) in the respective non-positive representative.

C. The result. Denote by $\mathcal{M}$ a category of torus bundles (of fixed dimension), endowed with homeomorphisms between the bundles; denote by $\mathcal{A}$ a category of the Cuntz-Krieger algebras $\mathcal{O}_A$ with $\det (A) = \pm 1$, endowed with stable isomorphisms between the algebras. Consider a (Cuntz-Krieger) map, $F : \mathcal{M} \rightarrow \mathcal{A}$, which acts by the formula $M_\alpha \mapsto \mathcal{O}_A$. The following is true.

Theorem 1 *The map F is a covariant functor, such that $H_1(M_\alpha; \mathbb{Z})$ and $\mathbb{Z} \oplus K_0(F(M_\alpha))$ are isomorphic abelian groups.*

The article is organized as follows. Theorem 1 is proved in section 2. There is no formal section on the preliminaries; all necessary results and concepts

are introduced in passing. (We encourage the reader to consult [1]–[5] for the details.) In section 3, an application of theorem 1 is considered.

2 Proof of theorem 1

The idea of proof consists in a reduction of the conjugacy problem for the automorphisms of T^n to the Cuntz-Krieger theorem on the flow equivalence of the subshifts of finite type (to be introduced in the next paragraph). There are no difficult parts in the proof, which is basically a series of observations. Moreover, theorem 1 follows from the results of P. M. Rodrigues and J. S. Ramos [3]. However, our accents are different and the proof is more direct (and shorter) than in the above cited work.

(i) The main reference to the subshifts of finite type (SFT) is [2]. A *full* Bernoulli n -shift is the set X_n of bi-infinite sequences $x = \{x_k\}$, where x_k is a symbol taken from a set S of cardinality n . The set X_n is endowed with the product topology, making X_n a Cantor set. The shift homeomorphism $\sigma_n : X_n \rightarrow X_n$ is given by the formula $\sigma_n(\dots x_{k-1}x_kx_{k+1}\dots) = (\dots x_kx_{k+1}x_{k+2}\dots)$. The homeomorphism defines a (discrete) dynamical system $\{X_n, \sigma_n\}$ given by the iterations of σ_n .

Let A be an $n \times n$ matrix, whose entries $a_{ij} := a(i, j)$ are 0 or 1. Consider a subset X_A of X_n consisting of the bi-infinite sequences, which satisfy the restriction $a(x_k, x_{k+1}) = 1$ for all $-\infty < k < \infty$. (It takes a moment to verify that X_A is indeed a subset of X_n and $X_A = X_n$, if and only if, all the entries of A are 1's.) By definition, $\sigma_A = \sigma_n | X_A$ and the pair $\{X_A, \sigma_A\}$ is called a *SFT*. A standard edge shift construction described in [2] allows to extend the notion of SFT to any matrix A with the non-negative entries.

It is well known that the SFT's $\{X_A, \sigma_A\}$ and $\{X_B, \sigma_B\}$ are topologically conjugate (as the dynamical systems), if and only if, the matrices A and B are *strong shift equivalent* (SSE), see [2] for the corresponding definition. The SSE of two matrices is a difficult algorithmic problem, which motivates the consideration of a weaker equivalence between the matrices called a *shift equivalence* (SE). Recall, that the matrices A and B are said to be shift equivalent (over $\mathbb{Z}^+$), when there exist non-negative matrices R and S and a positive integer k (a lag), satisfying the equations $AR = RB$, $BS = SA$, $A^k = RS$ and $SR = B^k$. Finally, the SFT's $\{X_A, \sigma_A\}$ and $\{X_B, \sigma_B\}$ (and the matrices A and B) are said to be *flow equivalent* (FE), if the suspension flows of the SFT's act on the topological spaces, which are homeomorphic

under a homeomorphism that respects the orientation of the orbits of the suspension flow. We shall use the following implications:

$$SSE \Rightarrow SE \Rightarrow FE. \quad (3)$$

(The first implication is rather classical, while for the second we refer the reader to [2], p.456.)

We further restrict to the SFT's given by the matrices with determinant ± 1 . In view of Corollary 2.13 of [5], the matrices A and B with $\det(A) = \pm 1$ and $\det(B) = \pm 1$ are SE (over $\mathbb{Z}^+$), if and only if, matrices A and B are similar in $GL_n(\mathbb{Z})$.

Let now α and α' be a pair of conjugate automorphisms of T^n . Since the corresponding matrices A and A' are similar in $GL_n(\mathbb{Z})$, one concludes that the SFT's $\{X_A, \sigma_A\}$ and $\{X_{A'}, \sigma_{A'}\}$ are SE. In particular, the SFT's $\{X_A, \sigma_A\}$ and $\{X_{A'}, \sigma_{A'}\}$ are FE.

One can now apply the known result due to Cuntz and Krieger; it says, that the C^* -algebra $\mathcal{O}_A \otimes \mathcal{K}$ is an invariant of the flow equivalence of the irreducible SFT's, see p. 252 of [1] and its proof in Sect. 4 of the same work. Thus, the map F sends the conjugate automorphisms of T^n into the stably isomorphic Cuntz-Krieger algebras, i.e. F is a functor.

Let us show that F is a covariant functor. Consider the following commutative diagram:

$$\begin{array}{ccc} A & \xrightarrow{\text{similarity}} & A' = BAB^{-1} \\ F \downarrow & & \downarrow F \\ \mathcal{O}_A & \xrightarrow[\text{isomorphism}]{\text{stable}} & \mathcal{O}_{BAB^{-1}}, \end{array}$$

where $A, B \in GL_n(\mathbb{Z})$ and $\mathcal{O}_A, \mathcal{O}_{BAB^{-1}} \in \mathcal{A}$. Let g_1, g_2 be the arrows (similarity of matrices) in the upper category and $F(g_1), F(g_2)$ the corresponding arrows (stable isomorphisms) in the lower category. In view of the diagram, we have the following identities:

$$\begin{aligned} F(g_1 g_2) &= \mathcal{O}_{B_2 B_1 A B_1^{-1} B_2^{-1}} = \mathcal{O}_{B_2 (B_1 A B_1^{-1}) B_2^{-1}} \\ &= \mathcal{O}_{B_2 A' B_2^{-1}} = F(g_1) F(g_2), \end{aligned} \quad (4)$$

where $F(g_1)(\mathcal{O}_A) = \mathcal{O}_{A'}$ and $F(g_2)(\mathcal{O}_{A'}) = \mathcal{O}_{A''}$. Thus, F does not reverse

the arrows and is, therefore, a covariant functor. The first statement of theorem 1 is proved.

(ii) Let M_α be a torus bundle with a monodromy, given by the matrix $A \in GL_n(\mathbb{Z})$. It can be calculated, e.g. using the Leray spectral sequence for the fiber bundles, that $H_1(M_\alpha; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}^n / (A - I)\mathbb{Z}^n$. Comparing this calculation with the K -theory of the Cuntz-Krieger algebra, one concludes that $H_1(M_\alpha; \mathbb{Z}) \cong \mathbb{Z} \oplus K_0(\mathcal{O}_A)$, where $\mathcal{O}_A = F(M_\alpha)$. The second statement of theorem 1 follows. $\square$

3 Examples

Consider the following (three-dimensional) torus bundles M_{α_i} ($i = 1, 2, 3$)¹:

$$\begin{aligned} A_1^n &= \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}, \quad n \in \mathbb{Z}, \quad K_0(\mathcal{O}_{A_1^n}) \cong \mathbb{Z} \oplus \mathbb{Z}_n, \\ A_2 &= \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}, \quad K_0(\mathcal{O}_{A_2}) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2, \\ A_3 &= \begin{pmatrix} 5 & 1 \\ 4 & 1 \end{pmatrix}, \quad K_0(\mathcal{O}_{A_3}) \cong \mathbb{Z}_4. \end{aligned}$$

(Here the group $K_0(\mathcal{O}_{A_i})$ was calculated using a reduction of the matrix to its Smith normal form [2].) Notice, that the Cuntz-Krieger invariant $K_0(\mathcal{O}_{A_1^n}) \cong \mathbb{Z} \oplus \mathbb{Z}_n$ is a complete topological invariant of the family of bundles $M_{\alpha_1^n}$; thus, such an invariant solves a classification problem for these bundles. For the bundles M_{α_2} and M_{α_3} the Alexander polynomial:

$$\Delta_{A_2}(t) = \Delta_{A_3}(t) = t^2 - 6t + 1. \quad (5)$$

Thus, the Alexander polynomial alone cannot distinguish between the bundles M_{α_2} and M_{α_3} ; however, since $K_0(\mathcal{O}_{A_2}) \not\cong K_0(\mathcal{O}_{A_3})$, theorem 1 says that the bundles M_{α_2} and M_{α_3} are topologically distinct.

Acknowledgments. The paper is dedicated to Prof. J. Cuntz on occasion of his sixtieth birthday. I am grateful to Prof. W. Krieger for a copy of [3] and exciting discussions.

¹It is interesting, that by Thurston's Geometrization Theorem, the bundle $M_{\alpha_1^n}$ is a nilmanifold for any n , while bundles M_{α_2} and M_{α_3} are the solvmanifolds [4].

References

- [1] J. Cuntz and W. Krieger, A class of C^* -algebras and topological Markov chains, *Invent. Math.* 56 (1980), 251-268.
- [2] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge Univ. Press, 1995.
- [3] P. M. Rodrigues and J. S. Ramos, Bowen-Franks groups as conjugacy invariants for T^n -automorphisms, *Aequationes Math.* 69 (2005), 231-249.
- [4] W. P. Thurston, Three dimensional manifolds, Kleinian groups and hyperbolic geometry, *Bull. Amer. Math. Soc.* 6 (1982), 357-381.
- [5] J. B. Wagoner, Strong shift equivalence theory and the shift equivalence problem, *Bull. Amer. Math. Soc.* 36 (1999), 271-296.

THE FIELDS INSTITUTE FOR MATHEMATICAL SCIENCES, TORONTO,
ON, CANADA, E-MAIL: igor.v.nikolaev@gmail.com

*Current address: 101-315 Holmwood Ave., Ottawa, ON, Canada, K1S
2R2*